Asymptotes 2



RaiseMyMarks.com

2020



# Vertical Asymptotes


Vertical asymptotes at x = c occurs when one of the following limits occur:

$$\lim_{x \to c^+} f(x) \text{ or } \lim_{x \to c^-} f(x)$$

does not exist. That is,

 $\lim_{x \to c^+} f(x) = +\infty$  $\lim_{x \to c^+} f(x) = -\infty$  $\lim_{x \to c^-} f(x) = +\infty$  $\lim_{x \to c^-} f(x) = -\infty$ 

For a general rational function  $f(x) = \frac{p(x)}{q(x)}$ , f has a vertical asymptote x = c if q(c) = 0 and  $p(c) \neq 0$ .





Let's consider and example.

### Example

Determine any vertical asymptotes of the function

$$f(x) = \frac{x}{x^2 + x - 2}$$

and describe the behavirous.

Solution:

$$f(x) = \frac{p(x)}{q(x)}$$

where p(x) = x and  $q(x) = x^2 + x - 2$ .

$$q(x) = x^{2} + x - 2 = (x + 2)(x - 1) \Rightarrow x = -2, 1$$

	-2		0	1
x	-	-	+	+
x+2		+	+	+
x - 1	-	-	-	+
f(x)	-	+	-	+

$$\lim_{x \to -2^{-}} f(x) = -\infty$$
$$\lim_{x \to -2^{+}} f(x) = +\infty$$
$$\lim_{x \to 1^{-}} f(x) = -\infty$$
$$\lim_{x \to x^{+}} f(x) = +\infty$$


## Horizontal Asymptotes

When the limit of the function is a finite number as  $x \to \infty$  or  $x \to -\infty$ . Or, if,

$$\lim_{x \to +\infty} f(x) = L \text{ or } \lim_{x \to -\infty} f(x) = L$$

then, we say the line y = L is a horizatonal asymptote of f(x).





### Example

Determine the equation of all asymptotes of the graph

$$f(x) = \frac{2x^2 + 3x - 1}{x + 1}$$

**Solution:** To find the vertical asymptotes, we need to consider when the function f(x) is undefined.  $p(x) = 2x^2 + 3x - 1$  and q(x) = x + 1 = 0. Therefore, x = -1 is a vertical asymptote. Notes that  $p(-1) = 2 - 3 - 1 = -2 \neq 0$ . To find the horizontal asymptotes,

$$\lim_{x \to \infty} f(x) = \lim_{x \to \infty} \frac{2x^2 + 3x - 1}{x + 1}$$
$$= \lim_{x \to \infty} \frac{x^2 \left(2 + \frac{3}{x} - \frac{1}{x^2}\right)}{x \left(1 + \frac{1}{x}\right)}$$
$$= \lim_{x \to \infty} 2x = \infty, \text{ and}$$
$$\lim_{x \to -\infty} f(x) = -\infty$$




Therefore, no horizontal asymptotes. In order to find any *oblique asymptotes*, we need to divide q(x) = (x + 1) into  $p(x) = 2x^2 + 3x - 1$ .

$$\begin{array}{r} 2x+1 \\ x+1 \overline{) \quad 2x^2+3x-1} \\ -(2x^2+2x) \\ \hline x -1 \\ -(x+1) \\ \hline -2 \end{array}$$

Therefore,

$$f(x) = \frac{2x^2 + 3x - 1}{x + 1} = 2x + 1 - \frac{2}{x + 1}$$
$$\lim_{x \to \infty} \frac{2}{x + 1} = 0 \quad , \quad \lim_{x \to -\infty} \frac{2}{x + 1} = 0$$
$$\therefore \text{ as } x \to \infty \quad , \quad f(x) \to 2x + 1 \text{ or}$$
$$\text{ as } x \to -\infty \quad , \quad f(x) \to 2x + 1.$$

Therefore, y = 2x + 1 is an oblique asymptote.





### Exercises

1. Check for discontinuities.

$$y = \frac{x}{x+5}$$

b)

c)

$$s = \frac{1}{(t-3)^2}$$

e)

d)

$$y = \frac{x^2 - x - 6}{x - 3}$$

 $f(x) = \frac{1}{e^x - 2}$ 

 $g(t) = \frac{3t^2 + 4}{t^2 - t}$ 

- 2. For each of the functions in #1 conduct a limit test to determine the behaviour on either side of the asymptote.
- 3. For each of the functions in #1 determine the horizontal and vertical asymptotes, if any.