

Increasing and Decreasing Functions

Raise My
MArks


RaiseMyMarks.com

2020

Increasing and Decreasing Functions

We say a function f is *decreasing* on an interval if for $x_1 < x_2$, $f(x_1) > f(x_2)$.



We say a function f is *increasing* on an interval if for any $x_1 < x_2$, $f(x_1) < f(x_2)$



If we consider the graph above, when $x < 0$ the function is decreasing. Notice that the tangents to the curve when it is decreasing are all negative or $f'(x) < 0$ for $x < 0$. When $x > 0$, the function is increasing. Notice that the tangents to the curve when $x > 0$ are all positive, $f'(x) > 0$, for $x > 0$. We can summarize this as follows:

For a continuous and differentiable function f ,
 f is increasing when $f'(x) > 0$ and f is decreasing when $f'(x) < 0$.

If we look at the graph below, we see that one graph has a maximum value and the other has a minimum value.



Notice in left figure we have a minimum value when $x = c$ or when $f'(c) = 0$. The point where this minimum occurs is $(c, f(c))$. In the right hand figure we have a similar situation except now we have a maximum value when $x = c$ or at point $(c, f(c))$. Again $f'(c) = 0$. This point $(c, f(c))$ is called a *critical point* of the function f .

Critical points

How do we find a critical point for a function f ? For a function $f(x)$

1. Find $f'(x)$.
2. Find the roots of $f'(x) = 0$. So, find the x values for which $f'(x) = 0$.
3. The x -values found in 2. can be labeled c . Evaluate $f(c)$ for each c value in 2.
4. $(c, f(c))$, for each c in 2. is a *critical point* for f .

How do we know if $f(c)$ is a relative maximum or relative minimum of the function? For a function f where $(c, f(c))$ is a critical point, if $f(c) < f(x)$ for x near c then $f(c)$ is a local minimum values. If $f(c) > f(x)$ for x near c then $f(c)$ is a local maximum value.

When graphing a function there are a few points and lines that can help us. The x and y intercepts; the critical points and the asymptotes. There are vertical and horizontal asymptotes.

Exercises

1. Find the points at which $f'(x) = 0$.

a) $f(x) = x^3 + 6x^2 + 1$

c) $f(x) = (2x - 1)^2(x^2 - 9)$

b) $f(x) = \sqrt{x^2 + 4}$

d) $f(x) = x^{2/3}(2x - 5)$

2. Find the critical points for the following functions.

a) $y = x^3 - 6x^2$

c) $y = \ln(x^2 - 3x + 4)$

b) $y = x^4 - 8x^2$