

Increasing and Decreasing Functions 2

**Raise My**  
**MArks**


RaiseMyMarks.com

2020

## Increasing and Decreasing Functions

We say a function  $f$  is *decreasing* on an interval if for  $x_1 < x_2$ ,  $f(x_1) > f(x_2)$ .

We say a function  $f$  is *increasing* on an interval if for any  $x_1 < x_2$ ,  $f(x_1) < f(x_2)$





If we consider the graph above, when  $x < 0$  the function is decreasing. Notice that the tangents to the curve when it is decreasing are all negative or  $f'(x) < 0$  for  $x < 0$ . When  $x > 0$ , the function is increasing. Notice that the tangents to the curve when  $x > 0$  are all positive,  $f'(x) > 0$ , for  $x > 0$ . We can summarize this as follows:

For a continuous and differentiable function  $f$ ,  
 $f$  is increasing when  $f'(x) > 0$  and  $f$  is decreasing when  $f'(x) < 0$ .

### Maximum or Minimum

If we look at the graph below, we see that one graph has a maximum value and the other has a minimum value.



Notice in left figure we have a minimum value when  $x = c$  or when  $f'(c) = 0$ . The point where this minimum occurs is  $(c, f(c))$ . In the right hand figure we have a similar situation except now we have a maximum value when  $x = c$  or at point  $(c, f(c))$ . Again  $f'(c) = 0$ . This point  $(c, f(c))$  is called a *critical point* of the function  $f$ .

## Critical points

How do we find a critical point for a function  $f$ ? For a function  $f(x)$

1. Find  $f'(x)$ .
2. Find the roots of  $f'(x) = 0$ . So, find the  $x$  values for which  $f'(x) = 0$ .
3. The  $x$ -values found in 2. can be labeled  $c$ . Evaluate  $f(c)$  for each  $c$  value in 2.
4.  $(c, f(c))$ , for each  $c$  in 2. is a *critical point* for  $f$ .

## Relative maximum or minimum?

How do we know if  $f(c)$  is a relative maximum or relative minimum of the function?

1. For a function  $f$  where  $(c, f(c))$  is a critical point, if  $f(c) < f(x)$  for  $x$  near  $c$  then  $f(c)$  is a local minimum values. If  $f(c) > f(x)$  for  $x$  near  $c$  then  $f(c)$  is a local maximum value.
2. Another way to determine if a critical point  $c$  give a maximum or minimum value is to consider the second derivative. If  $f''(c) > 0$  then  $c$  gives a minimum value, that is  $f(c)$  is a minimum; if  $f''(c) < 0$  then  $c$  gives a maximum value, that is  $f(c)$  is a maximum.

## Exercises

1. Find the critical points for the following functions.

a)  $y = x^3 - 6x^2$

e)  $h(x) = -6x^3 + 18x^2 + 3$

b)  $y = x^4 - 8x^2$

f)  $s = -t^2e^{-3t}$

c)  $y = \ln(x^2 - 3x + 4)$

g)  $g(t) = t^5 + t^3$

d)  $y = xe^{4x}$

2. For each function in #1 determine if the critical point is a local maximum or minimum or neither.