# Rules of Differentiation

# Raise My KS

RaiseMyMarks.com

2020



# Rules of differentiation

There are a number of rules when taking the derivative of a function. B

### Constant Function Rule

If 
$$f(x) = K$$
 then  $f'(x) = 0$ .

### The Power Rule

If  $f(x) = x^n$  then  $f'(x) = nx^{n-1}$  where n is a real number,  $n \in \mathbb{R}$ .

### Constant Multiple Rule

If 
$$f(x) = Kg(x)$$
 where K is a constant then  $f'(x) = Kg'(x)$ .

### The Sum Rule

If f(x) and g(x) are differentiable functions and F(x) = f(x) + g(x) then F'(x) = f'(x) + g'(x).

### Difference Rule

If 
$$F(x) = f(x) - g(x)$$
 then  $F'(x) = f'(x) - g'(x)$ 

### The Product Rule

If 
$$F(x) = f(x)g(x)$$
 then  $F'(x) = f'(x)g(x) + f(x)g'(x)$ 

### Power Rule


If 
$$f(x) = [g(x)]^n$$
 then  $f'(x) = n[g(x)]^{n-1}g'(x)$ , where  $n \in \mathbb{Z}$ ,  $n$  is an integer.

### The quotient Rule

If 
$$F(x) = \frac{f(x)}{g(x)}$$
 then  $F'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$ 

Before we look at the chain rule for differentiation let's' look at the composition of functions. The *composition of two or more functions* can be thought of as taking the function of a function or





The domain of one function is the range of the other function. Given two functions f and g the composite function  $f \circ g$  is defined by,

$$(f \circ g)(x) = f(g(x))$$

The chain rule considers the derivative of the composition of two functions.

### The Chain Rule

If f and g are differentiable functions and  $F(x) = f \circ g(x)$  then

$$F'(x) = f'(g(x))g'(x)$$

Using Leibniz Notation, If y = f(u), u = g(x) then,

$$\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx}.$$

Note: The power function rule is a special case of the chain rule where  $f(x) = x^n$  and given some function g(x), the derivative of  $F(x) = f \circ g(x) = [g(x)]^n$  is

$$f'(x) = n[g(x)]^{n-1}g'(x).$$



## Exercises

Use the rules of differentiation to differentiate the following functions.

a) 
$$f(x) = 3x^{2/3} - 6x^{1/3} + x^{-1/3}$$
 g)  $s = -2t^2 + 7t$ 

g) 
$$s = -2t^2 + 7t$$

b) 
$$f(x) = \sqrt{x} + 6\sqrt{x^3} + \sqrt{2}$$

h) 
$$s = (t - 3)^2$$

c) 
$$f(x) = \frac{1+\sqrt{x}}{x}$$

i) 
$$y = \frac{1}{x} (x^2 + \frac{1}{x})$$

d) 
$$f(x) = 9x^{-2} + 3\sqrt{x}$$

$$j) \ y = \frac{\sqrt{x} - 2}{\sqrt{x}}$$

e) 
$$f(x) = 20x^5 + 3\sqrt[3]{x} + 17$$

k) 
$$y = \frac{1}{x} \left( x^2 + \frac{1}{x} \right)$$

f) 
$$f(x) = \frac{1+\sqrt{x}}{x}$$



l) 
$$y = \frac{3}{x^2} - \frac{4}{x^3}$$

m) 
$$y = -x^2 + 3x + 4$$