Zeros or Roots of Polynomials - 2



RaiseMyMarks.com

2020



Zeros and Roots of Polynomials

The zeros or roots of a polynomial are the values of the independent variable usually x that result in the polynomial having a value of zero. These are the points on the graph of the polynomial that intersect the x-axis. Let's start by looking at *quadratics* and see how many possible zeros or roots a quadratic can have and what the graph of that quadratic will look like.

Quadratic

What is a quadratic? A quadratic is a polynomial with degree of 2. The general form of a quadratic is given by,

$$f(x) = ax^2 + bx + c$$
, where $a, b, c \in \mathbb{R}$

Let's start by considering a quadratic with 2 real roots or zeros.

Two real roots

Let's consider the following example of a quadratic with 2 real roots.

$$f(x) = a(x + 2)(x - 3)$$

zeros of f : -2, 3
roots of f(x) = 0 : -2, 3

What does the graph of this quadratic look like? Let's assume a > 0.





When a > 1 we have a thinner, stretched out curve; when a < 1 we have a flatter curve. When a < 0 then our curve opens downwards. Since we have two real roots, the curve crosses the x-axis twice.

One real root

Let's consider the following example of a quadratic with only 1 real root.

 $f(x) = a(x+2)^{2}$ zeros of f: -2roots of f(x) = 0: -2

What does the graph of this quadratic look like? Again, let's assume a > 0.



When a > 1 we have a thinner, stretched out curve; when a < 1 we have a flatter, compressed curve; when a < 0, our curve opens downwards. Since we have only one real root, the curve touches the x-axis once.

Cubic

What is a cubic? A cubic is a polynomial of degree 3. Recall that the degree of a polynomial is the value of the highest power in the polynomial. A general cubic is given by,

$$f(x) = ax^3 + bx^2 + cx + d$$



Three real roots

Let's consider the following example of a cubic with 3 real roots.

$$f(x) = a(x+2)(x-1)(x-3)$$

zeros of $f : -2, 1, 3$
roots of $f(x) = 0 : -2, 1, 3$

What does the graph of this type of cubic look like? Again, we'll assume a > 0 for our sketch.



Since we have 3 real roots, the graph crosses the x-axis 3 times and at the zeros or roots,-2, 1 and 3.

Two real roots

Let's consider the following example of a cubic with 2 real roots.

$$f(x) = a(x+2)(x-3)^2$$

zeros of $f: -2, 3$
roots of $f(x) = 0: -2, 3$

What does the graph of this cubic look like? Assume that a > 0.





Since we have 2 real roots, the graph crosses or touches the x-axis 2 times. Notice that the factor $(x-3)^2$ contributes the zero x = 3 but is also squared. This zero is the point at which the graph just touches the x-axis.

One real roots

Let's consider the following example of a cubic with 1 real root.

$$f(x) = a(x+2)^{3}$$

zeros of $f: -2$
roots of $f(x) = 0: -2$

What does the graph of this cubic look like? Assume a > 0. Recall the parent function $f(x) = x^3$? We know what the graph of this function looks like. The graph of $f(x) = a(x + 2)^3$ is this parent function transformed. What are the transformations? The first is the multiplication of a > 0 which is a stretch or compression depending on whether a > 1 or 0 < a < 1. The second transformation is the translation horizontally to the left 2 units. So the shape of this graph is similar to x^3 except that the "centre" is at x = -2 where the graph crosses the x-axis.





Here are some other examples of what a cubic may look like with one root



Quartic Four real roots

f(x) = a(x+2)(x-1)(x-3)(x+4)zeros of f: -2, 1, 3, -4 roots of f(x) = 0: -2, 1, 3, -4





Three real roots

$$f(x) = a(x+2)(x-1)(x-3)^{2}$$

zeros of $f: -2, 1, 3$
roots of $f(x) = 0: -2, 1, 3$





Two real roots



Note: For the above examples, the diagrams are just one representation of that particular case. There may be other diagrams representing each situation.

Exercises

- 1. Which functions are polynomial functions?
 - a) $f(x) = 3x^3 + 4x 2$ b) $f(x) = \frac{x^2 + 2x - 1}{x - 1}$ c) $f(x) = \sqrt{x^2 + 2x + 3}$ d) $f(x) = e^{2x - 2}$ e) $f(x) = -5x^5 + 3x - 1$ f) $f(x) = (x - 1)(x + 3)(x^2 - 2)^2$

2. Draw a sketch of the following functions.

- a) $y = (x-2)(x-3)(x+1)^2$ c) $y = -2(x+3)^2 4$
- b) $y = (x+2)^2$ d) y = (x-2)(x+3)



Zeros or Roots of Polynomials 2 - Exercises

- e) $y = x^3 2x^2 + 10x 2$ f) $y = x^4 + 3x - 2x^2 - 1$ g) y = 3(x - 9)(x + 1)h) y = 2(x - 6)(x + 4)i) $y = (x - 5)^2 + 6$
- j) $y = -\frac{1}{3}(x+4)^2$ k) $y = (x^2-4)^2$ l) $y = (x-1)x^2$ m) $y = (x-1)^3 + 4$