

Remainder Theorem

**Raise My**  
**Marks**

RaiseMyMarks.com

2021

## Remainder Theorem

The remainder theorem gives a relationship between the dividend  $f(x)$  and the remainder,  $r(x)$ .

### Example

$f(x) = x^3 - x^2 - 4x - 2$  divided by  $d(x) = x + 2$ . First let's consider,  $d(x) = x + 2 = 0$  and solve for  $x$ ,

**Solutions:**

$$d(x) = x + 2 = 0 \implies x = -2$$

Now, let's consider  $f(-2)$ .

$$\begin{aligned} f(-2) &= (-2)^3 - (-2)^2 - 4(-2) - 2 \\ &= -8 - 4 + 8 - 2 \\ &= -6 \end{aligned}$$

Now let's divide  $f(x) \div d(x)$  and see what we get.

$$\begin{array}{r} x^2 - 3x + 2 \\ x + 2 \overline{) x^3 - x^2 - 4x - 2} \\ \underline{-(x^3 + 2x^2)} \phantom{- 2} \\ -3x^2 - 4x \phantom{- 2} \\ \underline{-(-3x^2 - 6x)} \phantom{- 2} \\ 2x - 2 \phantom{- 2} \\ \underline{-(2x + 4)} \\ -6 \end{array}$$

We have  $q(x) = x^2 - 3x + 2$  and  $r(x) = -6 = f(-2)$ .

**Example**

Let  $f(x) = x^3 - 4x^2 + 5x - 1$  be divided by  $d(x) = x - 2$ .

**Solutions:**

$$\begin{array}{r}
 x^2 - 2x + 1 \\
 \hline
 x - 2 \ ) \ x^3 - 4x^2 + 5x - 1 \\
 \underline{-(x^3 - 2x^2)} \\
 -2x^2 + 5x \\
 \underline{-(-2x^2 + 4x)} \\
 -x - 1 \\
 \underline{-(-x - 2)} \\
 1
 \end{array}$$

and  $d(x) = x - 2 = 0$  gives,  $x = 2$ ,

$$f(2) = 2^3 - 4(2)^2 + 5(2) - 1 = 8 - 16 + 10 - 1 = 1 = r(x).$$

**Remainder Theorem**

What does the Remainder Theorem say?

If  $f(x)$  is divided by  $x - p$ , giving a quotient  $q(x)$  and a remainder  $r$  then  $r = f(p)$ .

**Example**

Find the remainder when  $f(x) = x^3 - 4x^2 + 5x - 1$  is divided by  $2x - 3$ .

Let's rewrite  $2x - 3$  in the form  $x - p$ ,

$$2x - 3 = 2 \left( x - \frac{3}{2} \right).$$

**Solutions:** Then by the remainder theorem,

$$\begin{aligned}f(x) &= f\left(\frac{3}{2}\right) = \left(\frac{3}{2}\right)^3 - 4\left(\frac{3}{2}\right)^2 + 5\left(\frac{3}{2}\right) - 1 \\&= \frac{27}{8} - \frac{36}{4} + \frac{15}{2} - 1 \\&= \frac{27}{8} - \frac{72}{8} + \frac{60}{8} - \frac{8}{8} \\&= \frac{7}{8} \\&= r\end{aligned}$$

### Example

What is the remainder when,  $x^3 - 4x^2 + 2x - 6$  is divided by  $x + 1$ ?

**Solutions:** So,

$$d(x) = x + 1 = 0 \implies x = -1 \text{ so, } p = -1.$$

$$\begin{aligned}r &= f(p) = f(-1) = (-1)^3 - 4(-1)^2 + 2(-1) - 6 \\&= -1 - 4 - 2 - 6 \\&= -13\end{aligned}$$

Therefore, the remainder is  $-13$ .

**Exercises**

1. Use the Remainder Theorem to find the remainder of the following:

(a)  $(x^3 - 4x^2 + 2x + 6) \div (2x + 3)$

(b)  $(3x^5 - 5x^2 + 4x + 1) \div (2x - 1)$

(c)  $(4x^3 + 9x - 10) \div (x - 1)$

(d)  $(6x^2 - 10x + 7) \div (3x + 1)$

(e)  $(x^4 - x^3 + x^2 - 3x + 4) \div (x - 5)$

2. Perform the following,

$$(a) (x^4 - 4x^3 + 3x^2 - 3) \div (x^2 + x - 2)$$

$$(b) (x^3 + 2x^2 - x - 2) \div (x - 1)$$

$$(c) (3x^3 + x + 2) \div (3x - 1)$$

$$(d) (6x^3 + 31x^2 + 25x - 12) \div (2x + 3)$$

$$(e) (4x^4 + 8x^3 - x^2 + x + 3) \div (x - 5)$$